

NeuroVue® Publications

- Wallace MM, Kavianpour SM, Gabriele MM. 2013. *Ephrin-B2 Reverse Signaling Is Required for Topography but Not Pattern Formation of Lateral Superior Inputs to the Inferior Colliculus*. **J. Comp. Neuro.**, 521:1585-1597.
- Kotak VC, Pendola LM, Rodriguez-Contreras A. 2012. *Spontaneous activity in the developing gerbil auditory cortex in vivo involves GABAergic transmission*. **Neuroscience** (Epub ahead of print).
- Duncan J, Kersigo J, Gray B and Fritzsich B. 2011. *Combining Lipophilic dye, in situ Hybridization, Immunohistochemistry and Histology*. **J. Visualized Experiments**, Issue 49, pub 3/17/2011 (ID: 2451).
- Kersigo J, D'Angelo A, Gray BD, Soukup GA, Fritzsich B. 2011. *The role of sensory organs and the forebrain for the development of the craniofacial shape as revealed by Foxg1-cre-mediated microRNA loss*. **Genesis**. Apr, 49 (4): 326-41
- Budefeld T, Tobet SA, Majdic G. 2011. *Altered position of cell bodies and fibers in the ventromedial region in SF-1*. **Exp. Neuro.** 232, 176-184.
- Maklad A, Kamel S, Wong E, Fritzsich B. 2010. *Development and organization of polarity-specific segregation of primary vestibular afferent fibers in mice*. **Cell Tissue Res** 340: 303-321.
- Fritzsich B, Dillard M, Lavado A, Harvey NL, Jahan I. 2010. *Canal cristae growth and fiber extension to the outer hair cells of the mouse ear require Prox1 activity*. **PLoS ONE**, 5, 2, e9377 (Feb).
- Tonniges J, Hansen M, Duncan J, Bassett MJ, Fritzsich B, Gray BD, Easwaran A, Nichols MG. 2010. *Photo- and bio-physical characterization of novel violet and near-infrared lipophilic fluorophores for neuronal tracing*. **J Microscopy**, 239 (2), 117-134.
- Pan, N., Jahan, I., Kersigo, J., Kopecky, B., Santi P., Johnson, S., Schmitz, H., Fritzsich, B. 2010. *Conditional deletion of Atoh1 using Pax2-Cre results in viable mice without differentiated cochlear hair cells that have lost most of the organ of Corti*. **Hearing Research**; 275, 66-80.
- Soukup GA, Fritzsich B, Pierce ML, Weston MD, Jahan I, et al. 2009. *Residual microRNA expression dictates the extent of inner ear development in conditional Dicer knockout mice*. **Dev Biol** 328: 328-341.
- Pan N, Jahan I, Lee JE, Fritzsich B. 2009. *Defects in the cerebella of conditional Neurod1 null mice correlate with effective Tg(Atoh1-cre) recombination and granule cell requirements for Neurod1 for differentiation*. **Cell Tissue Res** 337: 407-428.
- Maricich SM, Xia A, Mathes EL, Wang VY, Oghalai JS, Fritzsich B, Zoghbi HY. 2009. *Atoh1-lineal neurons are required for hearing and for the survival of neurons in the spinal ganglion and brainstem accessory auditory nuclei*. **J. Neurosci.**, sep 9; 29(36): 11123-33
- Nichols DH, Pauley S, Jahan I, Beisel KW, Millen KJ, Fritzsich B. 2008. *Lmx1a is required for segregation of sensory epithelia and normal ear histogenesis and morphogenesis*. **Cell Tissue Res** , 334 (3), 339-358

- Jensen-Smith H, Gray B, Muirhead K, Ohlsson-Wilhelm B, Fritzsche B. 2007. *Long-Distance Three-Color Neuronal Tracing in Fixed Tissue Using NeuroVue Dyes. Immunological Inv.* **36**, 5-6, 763-789.
- Kikkawa YS, Pawlowski KS. 2007. *Cochlear neuronal tracing for frequency mapping with Dil, NeuroVue, and Golgi methods. Acta Otolaryngol Suppl.* Dec; (559):19-23.
- Hsieh CY, Cramer KS. 2006. *Deafferentation Induces Novel Axonal Projections in the Auditory Brainstem After Hearing Onset. J Comp Neurol* 497: 589-599 NeuroVue Red was used for all figures except Figure 2D, for which both NeuroVue Red and Dil were used, and Figure 5A, for which Dil was used (K. Cramer, personal communication).
- Matei VA, Feng F, Pauley S, Beisel K, Nichols MG, Fritzsche B. 2006. *Near infrared laser illumination transforms the fluorescence absorbing X-Gal reaction product BCI into a transparent, yet brightly fluorescent substance. Brain Res Bull.* 70:33-43. NeuroVue Maroon (previously PTIR271; see Fritzsche, Dev. Dyn. 2005, below, for methods)
- Morris JK, Maklad A, Hansen LA, Feng F, Sorensen C, Lee KF, Macklin WB, Fritzsche B. 2006. *A disorganized innervation of the inner ear persists in the absence of ErbB2. Brain Res.* 1091: 186- 199 NeuroVue Maroon, NeuroVue Red
- Beisel, K.W., Wang-Lundberg, Y., Maklad, A. and Fritzsche, B. 2005. *Development and evolution of the vestibular sensory apparatus of the mammalian ear. J Vestibular Research,* 15:225-241.
- Fritzsche B, Muirhead KA, Feng F, Gray BD, Ohlsson-Wilhelm BM. 2005. *Diffusion and imaging properties of three new lipophilic tracers, NeuroVue Maroon, NeuroVue Red and NeuroVue Green and their use for double and triple labeling of neuronal profile. Brain Res Bull.,* 66:249-258.
- Fritzsche B, Matei VA, Nichols DH, Birmingham N, Jones K, Beisel KW, Wang VY. 2005. *Atoh1 null mutants show directed afferent fiber growth to undifferentiated ear sensory epithelia followed by incomplete fiber retention. Dev Dyn,* 233: 570-583. NeuroVue Maroon (previously PTIR271), NeuroVue Red (previously PTIR278)
- Fritzsche B, Pauley S, Matei V, Katz DM, Xiang M, Tessarollo L. 2005. *Mutant mice reveal the molecular and cellular basis for specific sensory connections to inner ear epithelia and primary nuclei of the brain. Hear Res,* 206: 52-63. NeuroVue Maroon (previously PTIR271).
- Fritzsche B, Jackson Lab Presentation. 2005. http://www.biomedsci.creighton.edu/facilities/nccb/media/Jackson_lab_presentation.ppt
- Matei V, Pauley S, Kaing S, Rowitch D, Beisel KW, Morris K, Jones K, Fritzsche B. 2005. *Smaller epithelia in Neurog1 null mice are related to earlier hair cell terminal mitosis. Dev Dyn,* 234:633-650. NeuroVue Maroon (previously PTIR271) was used for Figure 8 (B. Fritzsche, personal communication)
- Pauley, S., Matei, V., Beisel, K.W., Fritzsche, B. 2005. *Wiring the Ear to the Brain: the Molecular Basis of Neurosensory Development, Differentiation, and Survival. In: Development of the ear.* Eds. M. Kelley and D. Wu, **SHAR**, Springer Verlag, 85-120. NeuroVue Maroon (previouslyPTIR271) was used in Figure 4; see

- Tessarollo, J *Neurosci* 2004, below, for methods (B. Fritzsich, personal communication)
- de Caprona MD, Beisel KW, Nichols DH, Fritzsich B. 2004. *Partial behavioral compensation is revealed in balance tasked mutant mice lacking otoconia*. **Brain Res Bull** 64:289-301. Both NeuroVue Maroon (previously PTIR271) and NeuroVue Red (previously PTIR278) were used in Figure 8 (B. Fritzsich, personal communication).
 - Fritzsich B, Tessarollo L, Coppola E, Reichardt LF. 2004. *Neurotrophins in the ear: their roles in sensory neuron survival and fiber guidance*. **Prog Brain Res** 146:265-278. NeuroVue Maroon (formerly PTIR271) and Dil were used in Figure 2 (B. Fritzsich, personal communication)
 - Hellard D, Brosenitsch T, Fritzsich B, Katz DM. 2004. *Cranial sensory neuron development in the absence of brain-derived neurotrophic factor in BDNF/Bax double null mice*. **Dev Biol** 275:34-43. NeuroVue Maroon (previously PTIR271).
 - Gurung B, Fritzsich B. 2004. *Time course of embryonic midbrain and thalamic auditory connection development in mice as revealed by carbocyanine dye tracing*. **J Comp Neurol** 479:309-327. NeuroVue Maroon (previously PTIR271), and NeuroVue Red (previously PTIR278) used.
 - Maklad A, Fritzsich B, Hansen LA. 2004. *Innervation of the maxillary vibrissae in mice as revealed by anterograde and retrograde tract tracing*. **Cell. Tissue Res.**, 315: 167-180. NeuroVue Maroon was used for figure 6 (B. Fritzsich, personal communication)
 - Fritzsich B. 2003. *Development of inner ear afferent connections: forming primary neurons and connecting them to the developing sensory epithelia*. **Brain Res Bull.**, 60:423- 433. NeuroVue Maroon (previously PTIR271) and NeuroVue Green (previously PTIR281) were used in Figure 4a,4c,and 4d; NeuroVue Maroon and PKH26 were used for cover image (B. Fritzsich, personal communication)
 - Gu C, Rodriguez ER, Reimert DV, Shu T, Fritzsich B, Richards LJ, Kolodkin AL, Ginty DD. 2003. *Neuropilin-1 Conveys Semaphorin and VEGF Signaling during Neural and Cardiovascular Development*. **Dev Cell** 5:45-57. NeuroVue Maroon (formerly PTIR271) was used in Figure 3 (B. Fritzsich, personal communication)
 - Pauley S, Wright TJ, Pirvola U, Ornitz D, Beisel K, Fritzsich B. 2003. *Expression and function of FGF10 in mammalian inner ear development*. **Dev Dyn** 227:203-215. NeuroVue Maroon (formerly PTIR271) rather than Dil was used in Figure 7 (B. Fritzsich, personal communication)
 - Köbbert C, Apps R, Bechmann I, Lanciego JL, Mey J, Thanos S. 2000. *Current concepts of neuroanatomical tracing*. **Progress in Neurobiology** 62: 327-351.
 - Fritzsich, B, Nichols DH, Echelard Y, McMahon AP. 1995. *Development of midbrain and anterior hindbrain ocular motoneurons in normal and Wnt-1 knockout mice*. **J Neurobiol.** 27:457-469.
 - Fritzsich, B, Nichols DH, Echelard Y, McMahon AP. 1995. *Development of midbrain and anterior hindbrain ocular motoneurons in normal and Wnt-1 knockout mice*, **J Neurobiol.**, 27:457-469.